



Effects of Rhizobium, Nitrogen and Phosphorus Fertilizers on Growth, Nodulation, Yield and Yield Attributes of Soybean at Pawe Northwestern Ethiopia

Masresha Abitew Tarekegn^{1,*}, Kibebew Kibret²

¹Horticulture Department, College of Agriculture and Natural Resources, Debremarkos University, Debremarkos, Ethiopia

²School of Natural Resource Management and Environmental Science, College of Agriculture and Environmental Science, Haramaya University, Alemaya, Ethiopia

Email address: mass.ilt2005@gmail.com

Mobile: +251-910211292

ABSTRACT

Owing to the rising costs of chemical fertilizers and the growing environmental concerns, there is an ever increasing interest in the role of soil microorganisms in crop nutrition and soil fertility restoration. A field study was therefore conducted at Pawe Agricultural Research Center during 2010 main cropping season to determine the influence of *Bradyrhizobium japonicum* inoculation and N and P fertilizers application on nodulation, yield and yield attributes of soybean. Three levels of N (0, 11.5 and 23 kg N ha⁻¹); three levels of P (0, 23 and 46 kg P₂O₅ ha⁻¹) with two levels of Rhizobium were arranged in RCBD in factorial combinations with three replications. Nodule number, nodule fresh and dry weights, plant height, number of pods and seeds per plant, 100 seeds weight and grain yield responded significantly to the interaction effects of *B. japonicum* inoculation and application of N and P fertilizers but nodule volume, seed and biomass yields, and harvest index were significantly affected by the main effects of any one or more of the factors and interaction of any two of the factors. The maximum numbers of nodules of 80.26, fresh and dry weights of 3.77 and 0.99 gm./plant respectively; were measured by combined effect of 11.5 kg N/ha, 46 kg P₂O₅/ha and *B. japonicum*. Similarly the highest nodule volume of 3.53 and 3.27 ml/plant were measured after applications of 46 kg P₂O₅/ha with *B. japonicum* and with 11.5 kg N/ha respectively. The highest plant heights of 79.26 cm followed by 76.94 cm were measured by application of 46 kg P₂O₅/ha with *B. japonicum* and 46 kg P₂O₅/ha

itself respectively. Seed yield significantly increased to 11.91 gm /plant and 15.97 gm /plant following application of *B. japonicum* alone and 46 kg P₂O₅/ha with *B. japonicum* respectively. Applied 23 kg/ha brought the highest plant biomass of 27.25 gm/plant. Inoculation with 46 kg P₂O₅/ha and the combined effects of 11.5 kg N/ha, 46 kg P₂O₅/ha and *B. japonicum* resulted 73.93 and 80.66 number of pods per plant respectively. The maximum 100.60 numbers of seeds per plant were counted during inoculation of *B. japonicum* with 46 kg P₂O₅/ha. The maximum 100-seed weight of 16.96 gm and grain yield of 3151.88 kg/ha were produced by combined application of 11.5 kg N/ha, 46 kg P₂O₅/ha and *B. japonicum*. Each nodule attributes were significantly and positively correlated each other and with each yield and yield attributes. The results showed that growth and yield potential of soybean and an increase N₂ fixing can be achieved by using *B. japonicum* and P fertilizer with reduced level of N as starter fertilizer.

Keywords: *Bradyrhizobium japonicum*, Fertilizers, Nitrogen Fixation

1. INTRODUCTION

There is a general consensus on the need to address the problem of low soil fertility in a given region in order to improve agricultural productivity and thus food security. The augmentation of soil N is generally accomplished by many sources for supplying N to crops (Kovats *et al.*, 1985). Inorganic N fertilization is needed to alleviate its deficiency. However, it is costly and therefore out of reach of resource poor farmers. In addition, manure obtained from livestock could also be used as a cheap source of nutrients, but nutrient contents are often lower, which requires bulk application to satisfy plant nutrient demand (Kihanda, 1996). Biological nitrogen fixation (BNF) by leguminous crops can also supply sufficient N for optimum crop production (Laszlo and Jose, 2001). The contribution of the fixed N is a key factor in low input agricultural systems to sustain long-term soil fertility. Thus, the significance of BNF as the major mechanism for the recycling of N from the atmosphere to available forms in the biosphere need not be overemphasized.

Most agricultural soils of the tropics, including Ethiopia, are deficient in N and phosphorus (P) nutrients. These two nutrients often limit crop production in Ethiopia (Asgelil, 2000). Phosphorus deficiency followed by N is the major constraint in pulse production since it affects growth, nodule formation and development and N-fixation (Yohanes and Richer, 1999). Phosphorus has important effects on photosynthesis, N fixation, root development, flowering, seed formation, fruiting and improvement of crop quality (Brady, 2002). Symbiotic N fixation has a high P demand because the process consumes large amounts of energy (Schulze *et al.*, 2006) and energy generating metabolism strongly depends upon the availability of P (Plaxton, 2004). Therefore, N₂ fixation is very sensitive to P deficiency due to reduction in nodule mass and decreased ureide production (Sinclair and Vadez, 2002).

Similarly, mineral N fertilization is a crucial factor in oil seeded legume production (Rathke *et al.*, 2005). Even though BNF offers an alternative to the use of expensive ammonium based N fertilizer, the high yielding agricultural systems are difficult to sustain solely on BNF. So supplementation with mineral N might then be necessary for maximal yield of grain legumes (Zuberer, 1998). The effects of N and P fertilization on growth, yield and yield components and nodulation of legumes are not well investigated. Moreover, no studies have been conducted to find out the effects of N and P application on soybean growth, yield

and its nodulation in Pawe area soils. Therefore, there is a need to generate such information for improving productivity of soybean in the region. In view of this the present study was conducted with the following specific objectives;

- To examine the effect of *Bradyrhizobium japonicum* inoculation on growth, nodulation, yield and yield components of soybean,
- To evaluate the effects of N and P fertilizer application on the growth, nodulation, yield and yield components of soybean, and
- To examine the interactive effects of *Bradyrhizobium japonicum* with the application of N and P fertilizer.

2. MATERIALS AND METHODS

Description of Experimental Area

The experiment was conducted at Pawe the Agricultural Research Center (PARC) main station during the 2010 main cropping season. The PARC is found in Pawe district which is situated in Metekel Zone of Benishangul-Gumuz National Regional State in northwestern part of Ethiopia. Geographically, PARC is located at 11°09" N latitude and 36°03" E longitudes. It is 575 km in the Northwest direction of Addis Ababa. The area is characterized with an average annual rainfall of 1300 mm and monomodal rainfall distribution of intense storms that extend from May to October or November (Behailu, 2006). The mean annual maximum and minimum temperatures are 32 and 16 °C, respectively.




Figure 1. Monthly mean minimum and maximum temperatures and total rainfall of the experimental station (2010).

The trends of monthly mean minimum and maximum temperatures and rainfall of the 2010 cropping season are shown in Figure 1. From Figure 1, it can be seen that the rainfall increases sharply from the end of May to August. Maximum rainfall was recorded in July and August, which indicates that the crop was free of water deficit stress. But the rainfall then declined and ceased in October, which was at the physiological maturity of the crop. Therefore, the development or full growth period of the crop was not faced with any water deficit.

Experimental Treatments, Design and Procedures

The treatments were arranged in randomized complete block design (RCBD) in factorial combinations with three replications. The treatments were three levels of N fertilizer (0, 11.5 and 23 kg N ha⁻¹), three levels of P fertilizer (0, 23 and 46 kg P₂O₅ ha⁻¹) and two levels of *Bradyrhizobium japonicum* (TAL-379) inoculation (un-inoculated and inoculated). A total of 18 treatments were used in the experiment.

The size of each plot was 4 m × 2.4 m (9.6 m²). The space between plots was 1m and the space between the blocks was 2 m. Each plot contained 4 planting rows and the space between rows was 60 cm. At planting, two soybean seeds were seeded at 5 cm distance within a row and at 4 cm depth. Plant population was maintained by thinning at four to six leaf stages. The middle two rows were used for data collection and plant sampling for tissue analysis. Seeds were hand planted in rows on June 18. All standard cultural practices were applied throughout the growth period. Urea was used as N source while P was applied as triple super phosphate (TSP).

Bradyrhizobium japonicum Inoculant Preparation and Seed Inoculation

Seeds were treated with carrier based inoculants containing *Bradyrhizobium japonicum* at the rate of 10 g per kg of seed (Rice *et al.*, 2001). In order to ensure that the inoculum sticks to the seed, the required quantity of inoculants were suspended in 1:1 ratio in 10% sucrose solution so that the dry seeds were thoroughly mixed with the thick slurry of sugar solution (Subba Rao, 1988). After mixing, seeds were air-dried in the shade for 15 minutes and sown within an hour (Somasegaran and Hoben, 1985).

Soil Sampling and Preparation for Analysis

About 1 kg pre sowing surface soil sample was collected by means of auger from different spots of the experimental field at the depth of 0-30 cm and bulked together to get a representative composite soil sample based on the procedure outlined in Sahlemedhin and Taye (2000). Then, air-dried and crushed soil samples were thoroughly mixed and packed in a polythene bag, labeled and stored in the laboratory for analysis.

Chemical and Physical Analysis of Soil Properties

Soil samples were analyzed at the PARC Soil Laboratory. Soil particle size was analyzed using the Bouyoucos hydrometer method (Day, 1965) and textural class was identified on the basis of the USDA textural triangle. Soil pH was measured potentiometrically in 1:2.5 soil-water suspensions with standard glass electrode pH meter (van Reeuwijk, 1992). The Walkley and Black (1934) method was used to determine organic carbon content of the soil. The total N content of the soil was determined using the Kjeldhal digestion, distillation and titration

procedure (Jackson, 1958). Available P was analyzed based on the Bray II method (Bray and Kurtz, 1954). The ammonium acetate method was employed to determine the cation exchange capacity of the soil.

Determination of Shoot Nitrogen and Phosphorus Contents

Five plants were randomly sampled from the middle two rows of each plot at mid-flowering stage. The plant samples were then oven dried at 70 °C to a constant weight, and grounded to pass through 1 mm diameter sieve. Total N in shoot was estimated by the Kjeldhal digestion, distillation and titration method (Bremner and Mulvaney, 1982). Similarly, P content in plant tissue was estimated by wet digestion with a mixture of nitric acid (HNO₃) and perchloric acid (HClO₄). The P content was then determined by the vanadomolybdate yellow color using spectrophotometer at 460 nm (Islam *et al.*, 1992).

Data Collection

Five plants were randomly sampled from the two middle rows of each plot at mid-flowering stage of the plants. Then adhering soil particles were removed by washing the roots with their nodules gently with water over a metal sieve. Total number of nodules per plant was counted. The whole nodules from roots were picked; fresh and dry weight and volume of nodules per plant were recorded.

Five plants were randomly taken and plant height was measured with meter on tagged plants. The number of pods was determined at full maturity. Seed yield was harvested after picked up the pods from the randomly taken sampled plants. Five plants were again taken randomly and oven dried at 70 °C to constant weight to determine above ground biomass yield. Harvest index was calculated as the ratio of seed yield to above ground biomass yield. Samples of 100 seeds were taken from five randomly taken plants to determine 100 seed weight. Grain yield was recorded from plants harvested at physiological maturity. Grain yield was corrected for 10% moisture content using Draminski moisture meter and converted in to kilogram per hectare.

Statistical Analysis

The data were then subjected to analysis of variance for factorial experiment in RCBD design using Statistical Analysis System (SAS Version 9.0, 2004) Software. Mean separation was done using Fisher's least significant differences test (LSD) at 5% probability levels. Correlation analysis was also done between nodule characteristics, yield and yield attributes at correlation coefficients(r) value 0.05.

3. RESULTS AND DISCUSSION

Phosphorus and Nitrogen Contents of Tissue

Application of 11.5 and 23 kg ha⁻¹ N alone improved the tissue N content by 12.62 and 10.86%, respectively, over the control. This indicates that the use of high dose of starter N suppressed natural biological N fixation, which in turn might have resulted in less N content when artificial inoculation of *B. japonicum* were not made. On the other hand, the combined use of 11.5 and 23 kg ha⁻¹ N with *Bradyrhizobium japonicum* improved the tissue N content

by 17.82 and 28.27%, respectively compared to the control. Sarr *et al.* (2005) also previously reported significant increases in shoot N of soybean inoculated with *B. japonicum*.

Notwithstanding, remarkable improvements were observed when N and P fertilizers were combined with *B. japonicum*. The highest improvement of 34.77% was obtained when 11.5 kg ha⁻¹ N and 46 kg ha⁻¹ P₂O₅ were used in combination with *B. japonicum*. It is well known that *B. japonicum* inoculation and P increase *nitrogenase* activity and nodule mass that ultimately increases tissue N content, while addition of starter N fertilizer fulfills the immediate requirement of N of the plants at germination and these combinations lead to higher availability in soil and uptake of N by the plant.

Table 1. Interaction and main effects of *Bradyrhizobium japonicum* inoculation and mineral N and P fertilizers on plant tissue N and P contents of soybean at mid-flowering stage.

		Un-inoculated				Inoculated			
Applied N (kg ha ⁻¹)	Applied P (kg P ₂ O ₅ ha ⁻¹)				Applied P (kg P ₂ O ₅ ha ⁻¹)				
	0	23	46	Mean	0	23	46	Mean	
Plant tissue N content (%)									
0	3.233	3.655	3.416	3.435	3.514	3.795	3.851	3.72	
11.5	3.641	3.373	3.683	3.566	3.809	3.57	4.357	3.912	
23	3.584	3.697	3.556	3.612	4.147	3.879	4.006	4.011	
Mean	3.486	3.575	3.552	3.538	3.823	3.748	4.071	3.881	
Plant tissue P content (%)									
0	0.339	0.471	0.602	0.471	0.527	0.414	0.715	0.552	
11.5	0.471	0.565	0.527	0.521	0.339	0.301	0.339	0.326	
23	0.565	0.565	0.433	0.521	0.226	0.377	0.659	0.421	
Mean	0.458	0.534	0.521	0.504	0.364	0.364	0.571	0.433	

The response of P content in tissue with or without inoculation with *B. japonicum* was higher at the highest level of applied P (Table 1). At the P rate of 46 kg ha⁻¹ P₂O₅, inoculation with *B. japonicum* increased the P uptake by 18.78% compared to the un-inoculated one. Application of 23 and 46 kg ha⁻¹ P₂O₅ without inoculation of *B. japonicum* increased the P uptake by 38.94 and 77.58%, respectively. Furthermore, Olivera *et al.* (2004) also reported that phosphorus application to legumes increased plant biomass including nodule biomass and shoot P content due to the increased rate of N fixation.

Total P tissue content was further improved to a maximum of 110.9% over the control due to the application of 46 kg ha⁻¹ P₂O₅ with *B. japonicum* inoculation followed by 94%

through combined application of 23 kg ha⁻¹ N + 46 kg ha⁻¹ P₂O₅ and inoculation with *B. japonicum*. Increased P contents of straw, seed and P uptake in soybean due to combined application of P and *B. japonicum* inoculation was also reported by Moharram *et al.* (1994).

Nodulation Response of Soybean to Bradyrhizobium japonicum Inoculation and NP Fertilizers Application

Number of nodules per plant

The interaction of *Bradyrhizobium japonicum*, N and P had significant (P < 0.05) effect on the number of nodules. Application of 46 kg ha⁻¹ P₂O₅ and inoculation of *B. japonicum* alone significantly increased number of nodules per plant by 240.7% and 123%, respectively. However, application of N fertilizer alone did not bring about significant effect on the number of nodules (Table 2).

Table 2. Interaction effect of *Bradyrhizobium japonicum*, inoculation and mineral N and P fertilizers on nodule characteristics and plant height of soybean at Pawe.

		No. of nodules	F. weight (g)	D. weight (g)	Plant. Ht. (cm)
Un-inoculated					
N (kg ha ⁻¹) x P (P ₂ O ₅ kg ha ⁻¹)					
0	0	20.33 ^j	0.99 ^h	0.310 ⁱ	64.69 ^h
	23	32.93 ⁱ	2.049 ^{cd}	0.6 ^{gf}	74.2 ^c
	46	69.26 ^{bc}	2.33 ^c	0.8 ^{bc}	76.94 ^b
11.5	0	23.80 ^j	1.70 ^{def}	0.72 ^{de}	75.97 ^{bc}
	23	49.26 ^{fg}	1.37 ^{fgh}	0.65 ^{ef}	66.09 ^{gh}
	46	68.93 ^{bcd}	2.06 ^{cd}	0.82 ^{bc}	68.46 ^{ef}
23	0	23.66 ^j	1.65 ^{ef}	0.68 ^{de}	69.60 ^{ef}
	23	34.33 ^{hi}	1.03 ^h	0.44 ^h	65.46 ^h
	46	60.00 ^{de}	1.20 ^{gh}	0.45 ^h	67.74 ^{fg}
Inoculated					
0	0	45.40 ^g	1.98 ^{cde}	0.69 ^{de}	68.90 ^{ef}
	23	55.33 ^{ef}	1.56 ^{fg}	0.49 ^h	76.82 ^b
	46	73.13 ^{ab}	2.89 ^b	0.76 ^{bcd}	79.26 ^a
11.5	0	69.53 ^{bc}	2.28 ^c	0.74 ^{cd}	70.26 ^{de}

	23	63.33 ^{cde}	3.03 ^b	0.65 ^{ef}	71.96 ^d
	46	80.26 ^a	3.77 ^a	0.99 ^a	75.64 ^{bc}
23	0	42.06 ^{gh}	2.17 ^c	0.7 ^{de}	69.05 ^{ef}
	23	57.40 ^{ef}	1.42 ^{fg}	0.5 ^{gh}	68.36 ^{ef}
	46	71.66 ^{abc}	1.70 ^{def}	0.84 ^b	74.58 ^c
LSD (0.05)		9.11	0.38	0.08	2.16
CV (%)		10.53	11.95	7.53	1.83
SE (\pm)		2.61	0.10	0.02	0.61

Means followed by the same letter in a column are not significantly different at $P = 0.05$; No = Number, F = Fresh, D = Dry, Ht = height, SE = Standard error, CV = Coefficient of variation, LSD = Least significant difference.

Better nodule number per plant was observed due to the interaction of the three factors. Accordingly, inoculation of *B. japonicum* with combined application of 11.5 kg ha⁻¹ N and 46 kg ha⁻¹ P₂O₅, followed by combined use of *B. japonicum*, 0 kg ha⁻¹ N and 46 kg P₂O₅, and 23 kg ha⁻¹ N and 46 kg ha⁻¹ P₂O₅, significantly increased the mean number of nodules per plant as compared to most other treatments (Table 2). Greater number of nodules due to inoculation and N and P application suggested that there was better combining and symbiotic relationship between introduced *B. japonicum* and soybean. Wall *et al.* (2000) recognized that P in coincidence with the plant demand of N controls the nodule growth and alter the symbiotic processes between the legume and *Rhizobium*. This is in agreement with the findings of Revellin *et al.* (2000) and Abbasi *et al.* (2008), who reported that inoculation with *B. japonicum* significantly, increased the number of nodules per plant of soybean as compared to the control treatments.

Nodule volume per plant

The results of analysis of variance indicate that individual applications and the interaction effects of P by *B. japonicum* and N by P had significant effect ($P < 0.05$) on nodule volume per plant but not by the three factors. As a result, combined use of 11.5 kg ha⁻¹ N and 46 kg ha⁻¹ P₂O₅ increased mean nodule volume per plant significantly as compared to the other treatments. However, the combined application of 23 kg ha⁻¹ N with 46 kg ha⁻¹ P₂O₅ resulted in significantly lower mean nodule volume per plant than application of 46 kg ha⁻¹ P₂O₅ with 11.5 kg ha⁻¹ N. This indicates that the efficiency of soybean for nodule formation decreases with high level of N. High levels of inorganic N, especially nitrate N, have been shown to suppress nodulation of legumes (Herridge *et al.*, 1984).

However, the highest mean nodule volume per plant among the remaining treatments was observed when *B. japonicum* was used with 46 kg ha⁻¹ P₂O₅. This is because P is an essential ingredient for N-fixing bacteria and it has significant effects on the development of large number of nodules by providing energy required for N fixation.

Nodule fresh weight per plant

The results of analysis of variance indicate that all the three factors and their interactions affected fresh weight of nodules significantly ($P < 0.0001$). Soybean seed inoculation with *B. japonicum* resulted in greater mean nodule fresh weight than the control (Table 2). Like mean nodule number per plant, addition of either $11.5 \text{ kg ha}^{-1} \text{ N}$ or $23 \text{ kg ha}^{-1} \text{ N}$, showed insignificant change to the mean nodule fresh weight per plant. This can be again attributed to the inhibitory effect of N at higher levels on nodule number and size.

However, the maximum mean nodule fresh weight per plant was obtained by the interaction effect of *B. japonicum*, $11.5 \text{ kg ha}^{-1} \text{ N}$ and $46 \text{ kg ha}^{-1} \text{ P}_2\text{O}_5$ which was followed by after combined application of *B. japonicum*, $11.5 \text{ kg ha}^{-1} \text{ N}$ and $23 \text{ kg ha}^{-1} \text{ P}_2\text{O}_5$, and $0 \text{ kg ha}^{-1} \text{ N}$, $46 \text{ kg ha}^{-1} \text{ P}_2\text{O}_5$ and *B. japonicum* (Table 2). Nodule fresh weight was highly correlated with number of nodules indicating that nodule fresh weight increased as the number of nodules increased. Datsenko *et al.* (1997) concluded that number and weights of nodules are commonly used as the criteria of effective complementary interaction between macro and micro symbionts; thereby correlate on the whole with the rate of atmospheric nitrogen fixation. Apparently this was due to the formation of greater number of nodules from *B. japonicum* inoculated treatments. Jalaluddin (2005) also observed increment in fresh and dry weights of nodules in soybean in the *B. japonicum* inoculated plants as compared to control.

Nodule dry weight per plant

In line with the other nodulation parameters the nodule dry weight was also significantly affected ($P < 0.0001$) by the interaction of *Bradyrhizobium japonicum*, N and P. Soybean seed inoculation with *B. japonicum* and addition of $46 \text{ kg ha}^{-1} \text{ P}_2\text{O}_5$ increased the mean nodule dry weight per plant than the un-inoculated control (Table 2). However, there were no mean nodule dry weight differences among N fertilizer rates.

The maximum mean nodule dry weight per plant was obtained from the interaction effects of *B. japonicum*, $11.5 \text{ kg ha}^{-1} \text{ N}$ and $46 \text{ kg ha}^{-1} \text{ P}_2\text{O}_5$ which was followed by combined application of *B. japonicum*, $23 \text{ kg ha}^{-1} \text{ N}$ and $46 \text{ kg ha}^{-1} \text{ P}_2\text{O}_5$. This reveals that the combination of *B. japonicum* and P along with starter N produced the maximum nodule biomass. The result accords with the findings of Fatima *et al.* (2007) that using *B. japonicum* strains with P fertilizer increased the nodule dry weight as compared with the control. Presence of P and small N in the soil with *B. japonicum* strain might have individually or in combination positive effect on nodule weight and nodule volume parameters.

Soybean Growth and Biomass Response to Bradyrhizobium japonicum Inoculation and NP Fertilizers Application

Plant height

The results of analysis of variance indicate that the interaction of *Bradyrhizobium japonicum*, N and P was significant ($p < 0.0001$) on plant height. As a result, the use of 23 and $46 \text{ kg ha}^{-1} \text{ P}_2\text{O}_5$ alone significantly improved the mean plant height by 14.7 and 18.9%, respectively (Table 2). In consonance with the findings of this study, Abbasi *et al.* (2008), Mahmood *et al.* (2009) and Tahir *et al.* (2009) also reported an increase in plant height of soybean due to sole application of P. Sole application of 11.5 and $23 \text{ kg ha}^{-1} \text{ N}$ improved the mean plant height by 17.4 and 7.6%, respectively, over the control (Table 2). In line with this, a few studies showed that application of reduced amount of N as starter fertilizer could

improve growth by soybean (Ray *et al.*, 2006; Mrkovacki, *et al.*, 2008; Tahir *et al.*, 2009).

The *B. japonicum* improved the mean plant height of soybean by 6.5% than the un-inoculated plants. However, the maximum mean plant height improvement of 22.5 and 18.8% were obtained due to the interaction effects of *B. japonicum* and 46 kg ha⁻¹ P₂O₅, followed by combined application of *B. japonicum* and 23 kg ha⁻¹ N respectively, (Table 4). Significant improvement of 16.9% in mean plant height was obtained when 11.5 kg ha⁻¹ N, 46 kg ha⁻¹ P₂O₅ and *B. japonicum* were used. This was followed by a 15.3% increase due to the combined use of 23 kg ha⁻¹ N, 46 kg ha⁻¹ P₂O₅ and *B. japonicum* (Table 2). In general, the results indicate that combined use of N and P fertilizers with *B. japonicum* resulted in better mean plant height

Seed yield per plant

The results of analysis of variance indicate that the interaction of all the three factors did not significantly affect the seed yield, plant biomass and harvest index. However, the response of seed yield to seed inoculation of *B. japonicum*, application of starter N, and P and interactions of P by *B. japonicum* was significant. Accordingly, sole application of 11.5 kg ha⁻¹ N improved the mean seed yield per plant by 8% over the control but addition of 23 kg ha⁻¹ N alone showed the mean seed yield per plant statistically at par with the control (Table 3).

Seed inoculation of *B. japonicum* alone significantly increased mean seed yield per plant by 18.7% as compared to the un-inoculated control treatment. The findings of this study are also supported by the findings of Abbasi *et al.* (2008) who reported an increase in seed yield of soybean due to *B. japonicum* inoculation in Pakistan. The combined use of 46 kg ha⁻¹ P₂O₅ and *B. japonicum* significantly maximized the mean seed yield per plant by 59.2% as compared to the control. Furthermore, the combined use of 46 kg ha⁻¹ P₂O₅ and *B. japonicum* increased the mean seed yield of soybean by 34.1% over the use of seed inoculation of *B. japonicum* alone. This might be related to the increased nodulation through symbiosis between soybean and *B. japonicum*, which resulted in more N₂-fixation that leads to increased yield parameters.

Plant biomass per plant

Plant biomass was affected significantly by seed inoculation of *Bradyrhizobium japonicum* and sole application of N. As a result, soybean seed inoculation of *B. japonicum* alone increased the mean plant biomass per plant by 12.8% over the un-inoculated control. The result indicated that N fixation by *B. japonicum* enhanced the vegetative growth of soybean, which resulted in substantial increase in its biomass yield. Tahir *et al.* (2009) also reported increase in plant biomass due to inoculation of *B. japonicum* alone by 62.8% over the control. The maximum mean biomass yield per plant was obtained after application of 23 kg ha⁻¹ N. However, application of 11.5 kg ha⁻¹ N showed the mean biomass yield per plant statistically at par with the control (Table 3). The increase in biomass yield per plant was possibly because of supply of N with other soil mineral N form that was responsible for the highest vegetative growth of soybean. Mrkovacki *et al.* (2008) reported that maximum results for biomass yield were seen by applying 30 kg ha⁻¹ N to inoculated soybean instead of higher rates of N.

Harvest index per plant

Analysis of variance indicated that only independent application of N and P had significant variations on harvest index per plant. But inoculation and their interactions failed to respond significantly. The lowest mean harvest index per plant, even less than the control was recorded from the plots that received 23 kg ha⁻¹ N. Although application of 11.5 kg ha⁻¹ N showed statistically at par mean harvest index with the control, it resulted in 5.8% increase in mean harvest index per plant (Table 3). The decreased mean harvest index per plant with the increase of N fertilizer might be due to the influence of vegetative growth and increased above ground biomass yield, which reduced the harvest index.

Considering the main effects of P application on harvest index, the maximum mean harvest index per plant was obtained from application of 46 kg ha⁻¹ P₂O₅, which resulted in 19.1% increase over the control. The sole application of 23 kg ha⁻¹ P₂O₅, however, gave statistically similar mean harvest index per plant to that of the un-inoculated control treatment (Table 3). The increased mean harvest index per plant with the increase of P fertilizer rate might be due to the influence of greater fruit and seed setting than above ground biomass yield. The result found in this study is in agreement with the results of Malik *et al.* (2006) who reported that harvest index was significantly influenced by applied P in soybean crop.

Table 3. Main effects of *Bradyrhizobium japonicum*, inoculation and mineral N and P fertilizers on seed yield, plant biomass and harvest index of soybean at Pawe.

Treatments	Seed yield/plant (gm)	Plant biomass/plant (gm)	Harvest Index
Inoculation			
Un-inoculated	10.16 ^b	22.50 ^b	0.46
Inoculated	12.60 ^a	25.40 ^a	0.51
LSD (0.05)	1.13	2.3	NS
N (kg/ha)			
0	11.45 ^{ab}	22.22 ^b	0.52 ^a
11.5	12.36 ^a	22.38 ^b	0.55 ^a
23	10.32 ^b	27.25 ^a	0.38 ^b
LSD (0.05)	1.38	2.82	0.07
P (kg P ₂ O ₅ /ha)			
0	10.71 ^b	23.07	0.47 ^b
23	9.77 ^b	24.02	0.42 ^b
46	13.65 ^a	24.76	0.56 ^a

LSD (0.05)	1.38	NS	0.07
CV (%)	17.99	17.44	22.09
SE (\pm)	0.42	0.64	0.02

Means followed by the same letter in a column are not significantly different at $P = 0.05$; NS = No significance, SE = Standard error, CV = Coefficient of variation, LSD = Least significant difference.

Soybean Yield and Yield Attributes Response to Bradyrhizobium japonicum Inoculation and NP Fertilizers Application

Number of pods per plant

The results of analysis of variance indicated that individual application of *Bradyrhizobium japonicum*, N, P and their interactions had significant effect ($P < 0.0001$). Application of 23 and 46 kg ha⁻¹ P₂O₅ resulted in the same mean number of pods (Table 4). On the other hand, application of 11.5 kg ha⁻¹ N resulted in better mean pod number than the mean pod number obtained by application of 23 kg ha⁻¹ N and the control. Inoculation of *B. japonicum* by itself however showed equal mean pod number per plant than un-inoculated treatment (Table 4).

Also, combined application of 11.5 kg ha⁻¹ N and 46 kg ha⁻¹ P₂O₅ and 23 kg ha⁻¹ N and 46 kg ha⁻¹ P₂O₅ with no inoculation produced significantly greater number of mean pod number, improving it by 107.7 and 61.3%, respectively, (Table 4). Application of 23 and 46 kg ha⁻¹ P₂O₅ with *B. japonicum* resulted in 59.7 and 164.7% increase in mean pod number, respectively (Table 4). The significant improvements in mean pod number ranged from 33.2% to 110.7% due to application of 23 kg ha⁻¹ N and 11.5 kg ha⁻¹ N with *B. japonicum*, respectively. The maximum mean number of pods was recorded from plants that received 11.5 kg ha⁻¹ N and 46 kg ha⁻¹ P₂O₅ with *B. japonicum* inoculation (Table 4). An investigation conducted by Tahir *et al.* (2009) indicated that 94% increase of pod number per plant was recorded where 25 kg ha⁻¹ N was combined with P and *B. japonicum* on soybean in Pakistan.

Number of seeds per plant

The results of analysis of variance indicate that the interaction of *Bradyrhizobium japonicum*, N and P significantly ($P < 0.0001$) affected the number of seeds per plant. The minimum mean number of seeds per plant, even far less than that of the un-inoculated control, was found by the application of 23 kg ha⁻¹ N (Table 4). This could presumably be due to the fact that high rate of N fertilizer application significantly reduced number of nodules and nodule weights (Table 4) and the inhibitory effects of added N fertilizer to nodulation and N fixation lead to poor seed filling. The study by Cheema and Ahmad (2000) indicated that the dropping percentage in seeds number per plant with high N treatments ranged from 12-46% as compared to the control.

Bradyrhizobium japonicum inoculation alone gave better mean number of seeds count per plant than the control but when it was supplemented with 11.5 kg ha⁻¹ N and/or 46 kg ha⁻¹ P₂O₅, it increased the mean seed number per plant further (Table 4). This might be related to better N availability at the early growth stage of soybean, which in turn might have resulted in more nodule mass formation and adequate N fixation during growing season specifically seed setting

period; all of these contributing to the observed increased seed yield.

A considerable improvement in seed number per plant was also seen by independent applications of 46 kg ha⁻¹ P₂O₅ and 23 kg ha⁻¹ P₂O₅ as compared to the control (Table 4). However, the maximum mean number of seed count per plant was recorded by the combined applications of 11.5 kg ha⁻¹ N, 46 kg ha⁻¹ P₂O₅ and *B. japonicum* inoculation. The result confirms the findings of Tomar *et al.* (2004) who also observed significant increment in number of seeds per pod when *B. japonicum* inoculation was combined with different levels of P.

Hundred Seed weight

The results of analysis of variance indicate that the main effects of factors under consideration and their interactions showed significant differences on hundred seed weight except for the interaction of N by P. Seed inoculation resulted in higher mean hundred seed weight than un-inoculated treatment (Table 4). It increased hundred seed weight by 10.6% over the un-inoculated treatment. Increased in thousand seed weight resulted from *B. japonicum* inoculation was observed in soybean by Malik *et al.* (2006). Separate N applications contributed more or less the same output (Table 4). When applying 46 kg ha⁻¹ P₂O₅, reasonable hundred seed weight was measured than 23 kg ha⁻¹ P₂O₅. The maximum value of hundred seed weight was measured from mixed application of N and P inorganic fertilizers along with *B. japonicum* inoculation. It resulted in 26.9% increase in hundred seed weight followed by 20.8% through *B. japonicum* inoculation with 11.5 kg ha⁻¹ N + 46 kg ha⁻¹ P₂O₅ and 23 kg ha⁻¹ N + 46 kg ha⁻¹ P₂O₅, respectively (Table 4).

Table 4. Interaction effect of *Bradyrhizobium japonicum*, inoculation and mineral N and P fertilizers on yield and yield attributes of soybean at Pawe.

		Number of	Number of	100 seed	Grain
		pods/plant	seed/plant	Weigh (gm)	yield
					(kg/ha)
Un-inoculated					
N (kg ha ⁻¹) x P (P ₂ O ₅ kg ha ⁻¹)					
0	0	27.93 ^h	58.40 ^{gh}	13.36 ⁱ	1730.42 ^h
	23	70.13 ^b	83.46 ^c	13.85 ^{hi}	2511.25 ^{ef}
	46	68.66 ^b	93.13 ^b	16.02 ^{bcde}	2977.08 ^b
11.5	0	35.60 ^g	61.73 ^{efg}	15.34 ^{def}	2360.00 ^f
	23	29.40 ^h	70.53 ^d	13.74 ^{hi}	2602.50 ^{de}
	46	58.00 ^c	65.13 ^{def}	16.24 ^{ab}	3060.42 ^{ab}
23	0	32.80 ^h	54.60 ^h	14.8 ^{fg}	2171.46 ^g
	23	27.40 ^h	60.13 ^{fgh}	13.67 ^{hi}	2396.04 ^f

	46	45.06 ^{de}	60.86 ^{fg}	14.40 ^{gh}	2346.04 ^f
		Inoculated			
0	0	32.46 ^{gh}	78.13 ^c	14.78 ^{fg}	2638.33 ^{de}
	23	44.60 ^{de}	67.40 ^{de}	15.14 ^{fg}	2644.38 ^{de}
	46	73.93 ^b	100.60 ^a	16.11 ^{bcd}	3057.92 ^{ab}
11.5	0	58.86 ^c	91.33 ^b	15.29 ^{ef}	2763.75 ^{cd}
	23	48.93 ^d	61.13 ^{fg}	15.38 ^{cdef}	2910.21 ^{bc}
	46	80.66 ^a	120.00 ^a	16.96 ^a	3151.88 ^a
23	0	37.20 ^{fg}	58.26 ^{gh}	15.03 ^{fg}	2608.33 ^{de}
	23	42.80 ^{ef}	61.26 ^{efg}	15.26 ^{ef}	2622.71 ^{de}
	46	73.13 ^b	100.00 ^b	16.15 ^{bc}	2893.96 ^{bc}
LSD (0.05)		6.05	6.18	0.78	170.83
CV (%)		7.42	5.37	3.15	3.91
SE (\pm)		2.42	1.88	0.14	49.07

Means followed by the same letter in a column are not significantly different at $p = 0.05$; SE = standard error, CV = coefficient of variation, LSD = least significant difference.

Grain yield per hectare

The results of analysis of variance showed that the response of grain yield to the interaction effects of *Bradyrhizobium japonicum*, N and P was significant ($P < 0.0001$). Soybean seed inoculation with *B. japonicum* produced higher mean grain yield than that of the un-inoculated control (Table 4). The increase in grain yield could be attributed to increase in yield components of the crop through *B. japonicum* inoculated plots. Egamberdiyeva *et al.* (2004) reported increase in grain yield due to *B. japonicum* inoculation of soybean in Uzbekistan. Great grain yield variation was also observed among the P fertilizer rates. Addition of 46 kg ha⁻¹ P₂O₅ alone could yield more grain yield than 23 kg ha⁻¹ P₂O₅. However, better grain yield was harvested when 46 kg ha⁻¹ P₂O₅ was added with 11.5 kg ha⁻¹ N.

The final grain yield of a crop is a function of cumulative contribution of its various growth and yield parameters which are influenced by various agronomic practices. The highest grain yield was recorded when the three factors interacted to each other. As a result interaction of 11.5 kg ha⁻¹ N and 46 kg ha⁻¹ P₂O₅ along with *B. japonicum* inoculation gave the maximum grain yield. This was justified with the finding of Dubey (1998) who obtained highest grain yield of soybean when the plant was inoculated with *B. japonicum* in combination with N and P fertilizers.

Correlation Analysis of Nodulation and Yield Parameters

Correlation coefficient values exhibited that nodule number, volume, fresh and dry weights were highly significantly and positively correlated with each other. All the nodule characteristics were also correlated significantly and positively with each yield and yield attributes except to the biomass yield per plant. This indicates that the development of effective and promising nodules of the crop due to P supply and *B. japonicum* inoculation could promote N uptake through the process of BNF which ultimately enhances the final grain yield and yield attributes of the crop. The yield of plant is a dependent variable, depends upon all other growth and yield contributing components. Therefore, it is generally correlated with all other components.

4. SUMMARY AND CONCLUSIONS

In general, the response of P content in tissue with or without inoculation with *B. japonicum* was higher at the higher levels of applied P. Similarly, all the *B. japonicum* treatments used alone or in combination with inorganic fertilizers improved the tissue N content of soybean. Generally, the results of this study revealed that seed inoculation of *B. japonicum* and P application alone or in combination with *B. japonicum*, or P with small dose of N fertilizer could improve nodulation potential of soybean. Higher dose of N applied alone or in combination either with P or *B. japonicum* reduced the efficiency of *B. japonicum* for various nodule characteristics, growth parameters and yield traits of soybean. In general, starter N of 11.5 kg ha⁻¹ N with *B. japonicum* and 46 kg ha⁻¹ P₂O₅ resulted in better nodulation, growth and development, yield and yield attributes of soybean crop.

The results obtained in this work might have potential applications for increasing the productivity of soybean and enriching the soil with N. However, since the experiment is conducted for one season and at one location, it is difficult to give comprehensive recommendation on best combinations of inorganic fertilizers and *B. japonicum*. It is, therefore, necessary to repeat the experiment under various soil conditions and fertilizers rates with an appropriate symbiont strain. Therefore, it would be worthwhile to conduct a similar study in N depleted fields prevalent in smallholder production systems.

Acknowledgements

The authors would like to thank the staff members of the Pawe Agricultural Research Centre (PARC), for their material support, assistance, patience and their endless contributions for the success of this study. Mr. Musefa Redi and Mrs. Sebele, who assisted him in analyzing the soil and plant samples, deserve special gratitude. The authors would like also acknowledge the financial support of Ethiopian Ministry of Education and the Mizan-Tepi University for funding this research and sponsoring this training.

References

- [1] Abbasi, M. K., A. Majeed., A. Sadiq, and S. R. Khan, 2008. Application of Bradyrhizobium japonicum and phosphorus fertilization improved growth, yield and nodulation of soybean in the sub-humid hilly region of Azad Jammu and Kashmir,

- Pakistan. *Pak. J. Plant Prod. Sci.* 58, 368-376.
- [2] Asgelil Dibabe, 2000. Effect of fertilizer on the yield and nodulation pattern of faba bean on a Nitosol of Adet, North Western Ethiopia. *Ethio. J. Nat. Res.* 2, 237-244.
- [3] Behailu Kebede, 2006. Land cover, land use changes and agroforestry practices at Pawe resettlement district, northwestern Ethiopia. An MSc Thesis, Hawassa University, Wondo Genet College of Forestry and Natural Resources, Ethiopia.
- [4] Brady, N. C., 2002. Phosphorus and potassium. In: The nature and properties of soils. Prentice Hall of India, Delhi. 352 p.
- [5] Bray R.H. and L. T. Kurtz, 1954. Determination of total organic and available forms of phosphorous in soil. *J. Soil Sci.* 59, 39-45.
- [6] Bremner, J. M. and C. S. Mulvaney, 1982. Total nitrogen. pp. 595-624. In: A.L Page (eds.). Method of Soil Analysis. Agron. No. 9. Amer. Soc. Agron. Madison, WI, USA.
- [7] Cheema, Z. A. and A. Ahmad, 2000. Effects of urea on the nitrogen fixing capacity and growth of grain legumes. *Int. J. Agric. Biol.* 2(4), 388-394.
- [8] Datsenko, V. K., S. K. Laguta, E. P. Starchenkov, A. F. Antipchuk, and V. N. Rangelova, 1997. Efficiency of legume-rhizobia symbiosis in various soybean varieties and Bradyrhizobium japonicum cultures. *Fiziol. Biokhim. Kul't. Rast.* 29, 299-303.
- [9] Day, P. R., 1965. Hydrometer method of particle size analysis. pp. 562-563. In: C.A. Black (eds.). Methods of Soil Analysis. Agronomy Part II, No. 9. Amer. Soc. Agron. Madison, Wisconsin, USA.
- [10] Dubey, S. K., 1998. Response of soybean to biofertilizers with and without nitrogen, phosphorus and potassium on swell-shrink soil. *Ind. J. Agron.* 43, 546-549.
- [11] Egamberdiyeva, D., D. Qarshieva, K. Davranov, 2004. The use of Bradyrhizobium to enhance growth and yield of soybean in calcareous soil in Uzbekistan. *J. Plant Growth Regul.* 23, 54-57.
- [12] Fatima, Z., M. Zia, M.F. Chaudhary, 2007. Interactive effect of Rhizobium strains and phosphorus on soybean yield, nitrogen fixation and soil fertility. *Pak. J. Bot.* 39, 255-264.
- [13] Herridge, D. F., R. J. Roughley and J. Brockwell, 1984. Effects of Rhizobium and soil nitrate on establishment and functioning of the soybean symbiosis in the field. *Austr. J. Agric. Res.* 35, 146-161.
- [14] Islam, A. K. M. S., G. Kerven and J. Oweczkin, 1992. Methods of Plant Analysis. ACIAR 904 IBSRAM QC.
- [15] Jackson, M. L., 1958. Soil Chemical Analysis. Prentice Hall, Inc., Englewood Cliffs, New Jersey. Pp. 183-204.
- [16] Jalaluddin, M., 2005. Effect of inoculation with vsm-fungi and Bradyrhizobium on growth and yield of soybean in Sindh. *Pak. J. Bot.* 37: 169-173.
- [17] Kihanda, F. M., 1996. The role of farmyard manure in improvement of maize production in the sub-humid highlands of central Kenya. Ph.D Thesis, Reading

University, United Kingdom.

- [18] Kovats, A., L. Marton and L. Szabo, 1985. Analysis of the relation between humus and pH on the ground of results of soil investigations on farm-scale plots. *Plant Prod.* 34, 507-512.
- [19] Laszlo, M. and E. M. Jose, 2001. Effects of *Erotalaria juncea* L. and *Erotalaria spectabilis* ROTH on soil fertility and soil conservation in Hungary. *Acta Agronomica Óváriensis* 43, 1-8.
- [20] Mahmood, J., Y. A. Abayomi, M. O. Aduloju, 2009. Comparative growth and yield responses of soybean genotypes to phosphorus fertilizer application. *Afr. J. Biotech.* 8, 1030-1036.
- [21] Malik, M. A., M. A. Cheema, H. Z. Khan and M. A. Wahid, 2006. Growth and yield response of soybean (*Glycine max* L.) to seed inoculation and varying phosphorus levels. *J. Agric. Res.* 44(1), 47-53
- [22] Moharram, T. M. M., M. S. A. Safwat, M. M. Farghaly, 1994. Effect of inoculation rates and phosphorus fertilization on nitrogen fixation in soybean. *Afr. J. Crop Sci.* 2, 125-129.
- [23] Mrkovacki, N., J. Marinkovic, R. Rilmovic, 2008. Effect of n fertilizer application on growth and yield of inoculated soybean. *Not. Bot. Hort. Agrobot. Cluj.* 36, 48-51.
- [24] Olivera, M., N. Tejera, C. Iribarne, A. Ocana, C. Lluch, 2004. Growth, nitrogen fixation and ammonium assimilation in common bean (*Phaseolus vulgaris*). *Plant Physiol.* 121, 498-505.
- [25] Plaxton, W. C., 2004. Plant response to stress, biochemical adaptations to phosphate deficiency. Pp. 976-980. In: R. Goodman (ed.). *Encyclopedia of Plant and Crop Science*. Marcel Dekker, New York.
- [26] Rathke, G. W., O. Christen and W. Diepenbrock, 2005. Effects of nitrogen source and rate on productivity and quality of winter oilseed rape (*Brassica napus* L.) grown in different crop rotations. *Field Crop Res.* 94, 103-113.
- [27] Ray, J. D., L. G. Heatherly, F. B. Fritschi, 2006. Influence of large amounts of nitrogen applied at planting on non-irrigated and irrigated soybean. *Crop Sci.* 46, 52-60.
- [28] Revellin, C., G. Meunier, J. J. Giraud, G. Sommer, P. Wadoux, G. Catroux, 2000. Changes in the physiological and agricultural characteristics of peat-based *Bradyrhizobium japonicum* inoculants after long term storage. *Appl. Microbiol. Biotech.* 54, 206-211.
- [29] Rice, W. A., G. W. Clayton, N. Z. Lupwayi and P. E. Olsen, 2001. Evaluation of coated seeds as a *Rhizobium* delivery system for field pea. *Can. J. Plant Science* 81, 248-249.
- [30] Sahlemedhin Sertsue and Taye Bekele, 2000. Procedures for soil and plant analysis. National Soil Research Center, Ethiopian Agricultural Research Organization, Addis Ababa, Ethiopia. 110 p.
- [31] Sarr A., B. Diop, R. Peltier, M. Neyra, D. Lesueur, 2005. Effect of rhizobial inoculation methods and host plant provenances on nodulation and growth of *Acacia senegal* and

- Acacia nilotica*. *New Forests* 29, 75- 87.
- [32] SAS (Statistical Analysis System) Institute, 2004. The SAS system for windows, version 9.0. SAS Institute Inc., Cary, NC., USA.
- [33] Schulze, J., G. Temple, S. J. Temple, H. Beschow and C. P. Vance, 2006. Nitrogen fixation by white lupin under phosphorus deficiency. *Ann. Bot.* 98, 731-740.
- [34] Sinclair, T. R. and V. Vadez, 2002. Physiological traits for crop yield improvement in low N and P environments. *Plant and Soil.* 245, 1-15.
- [35] Somasegaran and Hoben, 1985. Methods in legume-Rhizobium technology. University of Hawaii Niftal Project and MIRCEM. Department of Agronomy and Soil Science, University of Hawaii.
- [36] Subba Rao, N. S., 1988. Rhizobium inoculants: Biofertilizer in agriculture and forestry, second ed. Oxford and IBH Publishing Co. Pvt Ltd, New Delhi, India. Pp. 16-76.
- [37] Tahir, M. M., M. K. Abbasi, N. Rahim, A. Khaliq and M. H. Kazmi, 2009. Effects of Rhizobium inoculation and NP fertilization on growth, yield and nodulation of soybean (*Glycine max* L.) in the sub-humid hilly region of Rawalakot Azad Jammu and Kashmir, Pakistan. *Afr. J. Biotech.* 8, 6191-6200.
- [38] Tomar, S. S., R. Singh and P. S. Singh, 2004. Response of phosphorus, sulphur and Rhizobium inoculation on growth, yield and quality of soybean. *Prog. Agric.* 4, 72-73.
- [39] Van Reeuwijk, L. P., 1992. Procedure for soil analysis, 4th ed, international soil reference and information center (ISRIC), Wageningen, Netherlands.
- [40] Walkley, A. and C. A. Black, 1934. An examination of different methods for determining soil organic matter and the proposed modification by the chromic acid titration method. *Soil Sci.* 37, 29-38.
- [41] Wall, L. G., A. Hellesten, K. Huss-Danell, 2000. Nitrogen, phosphorus and the ratio between them affect nodulation in *Alnus incana* and *Wfolium prattense*. *Symbiosis* 29, 91-105.
- [42] Yohanes Uloro and Richer, 1999. Phosphorus efficiency of different variety of *Phaseolus vulgaris* and *Sorgum bicolor* (L). Moend on an Altisols in Eastern Ethiopia Highlands. *Ethio. J. Nat. Res.* 1, 187-200.
- [43] Zuberer, D. A., 1998. Biological di-nitrogen fixation: Introduction and Non-symbiotic. In: Principles and Applications of Soil Microbiology. Prentice Hall, Inc. Simon & Schuster. A Viacom Company, Upper Saddle River, New Jersey, America.

(Received 02 February 2017; accepted 14 February 2017)